

ప్రొఫెసర్ జయశంకర్ తెలంగాణ వ్యవసాయ విశ్వవిద్యాలయం  
వ్యవసాయ వాతావరణ పరిశోధన కేంద్రం  
వ్యవసాయ పరిశోధన సంస్థ, రాజేంద్రనగర్, హైదరాబాద్.



బులెటిన్ నెం. XXXVI/54/2025

రాష్ట్ర రైతాంగానికి

వాతావరణ ఆధారిత వ్యవసాయ సలహాలు

తేది: 09.07.2025 (బుధవారం) నుండి 13.07.2025 (ఆదివారం) వరకు



గ్రామీణ కృషి మౌసమ్ సేవ పథకం (GKMS Project)

భారత వాతావరణ శాఖ, న్యూఢిల్లీ.

### గత మూడు రోజుల వాతావరణం:

గడిచిన మూడు రోజులలో రాష్ట్రంలో చాలా చోట్ల తేలికపాటి నుండి ఓ మోస్తరు వర్షాలు కురిసాయి. పగటి ఉష్ణోగ్రతలు 29 నుండి 35 డిగ్రీల సెల్సియస్ మరియు రాత్రి ఉష్ణోగ్రతలు 20 నుండి 26 డిగ్రీల సెల్సియస్ గా నమోదయ్యాయి.

### రాబోవు ఐదు రోజుల వాతావరణ విశ్లేషణ (ఈరోజు మధ్యాహ్నం 1300 గంటల ఆధారంగా):

హైదరాబాద్ వాతావరణ కేంద్రం వారు అందించిన సమాచారం ప్రకారం రాబోవు ఐదు రోజులలో కొన్ని చోట్ల తేలికపాటి నుండి ఓ మోస్తరు వర్షాలు కురిసే సూచనలున్నాయి. పగటి ఉష్ణోగ్రతలు 26 నుండి 36 డిగ్రీల సెల్సియస్ మధ్య మరియు రాత్రి ఉష్ణోగ్రతలు 21 నుండి 28 డిగ్రీల సెల్సియస్ మధ్య నమోదుకావచ్చు.

### హెచ్చరిక:

#### మొదటి రోజు:

ఉరుములు, మెరుపులు మరియు ఈదురుగాలులతో (గంటకు 40-50 కి.మీ.) కూడిన భారీ నుండి అతి భారీ వర్షాలు తెలంగాణ రాష్ట్రంలోని ఆదిలాబాద్, కొమరంభీం ఆసిఫాబాద్, నిర్మల్ జిల్లాలలో అక్కడక్కడ కురిసే అవకాశం ఉంది.

ఉరుములు, మెరుపులు మరియు ఈదురు గాలులతో (గంటకు 30-40 కి.మీ.) కూడిన భారీ వర్షాలు తెలంగాణ రాష్ట్రంలోని మంచిర్యాల, నిజామాబాద్ జిల్లాలలో అక్కడక్కడ కురిసే అవకాశం ఉంది.

ఉరుములు, మెరుపులతో కూడిన వర్షాలు తెలంగాణ రాష్ట్రంలోని జగిత్యాల, రాజన్న సిరిసిల్ల, కరీంనగర్, పెద్దపల్లి, జయశంకర్ భూపాలపల్లి, ములుగు, భద్రాద్రి కొత్తగూడెం, ఖమ్మం, నల్గొండ, సూర్యాపేట, మహబూబాబాద్, వరంగల్, హన్మకొండ, జనగాం, సిద్దిపేట, యాదాద్రి భువనగిరి, రంగారెడ్డి, హైదరాబాద్, మేడ్చల్ మల్కాజ్ గిరి, వికారాబాద్, సంగారెడ్డి, మెదక్, కామారెడ్డి జిల్లాలలో అక్కడక్కడ కురిసే అవకాశం ఉంది.

బలమైన ఈదురు గాలులు (గాలి వేగం గంటకు 30-40 కి.మీ.)తో తెలంగాణ రాష్ట్రంలోని అన్ని జిల్లాలలో అక్కడక్కడ వీచే అవకాశం ఉంది.

#### రెండవ రోజు:

ఉరుములు, మెరుపులు మరియు ఈదురు గాలులతో (గంటకు 30-40 కి.మీ.) కూడిన భారీ వర్షాలు తెలంగాణ రాష్ట్రంలోని ఆదిలాబాద్, కొమరంభీం ఆసిఫాబాద్, మంచిర్యాల, నిర్మల్ జిల్లాలలో అక్కడక్కడ కురిసే అవకాశం ఉంది.

ఉరుములు, మెరుపులతో కూడిన వర్షాలు తెలంగాణ రాష్ట్రంలోని నిజామాబాద్, జగిత్యాల, రాజన్న సిరిసిల్ల, కరీంనగర్, పెద్దపల్లి, జయశంకర్ భూపాలపల్లి, ములుగు, భద్రాద్రి కొత్తగూడెం, ఖమ్మం, నల్గొండ, సూర్యాపేట, మహబూబాబాద్, రంగారెడ్డి, హైదరాబాద్, మేడ్చల్ మల్కాజ్ గిరి, వికారాబాద్, సంగారెడ్డి, మెదక్, కామారెడ్డి జిల్లాలలో అక్కడక్కడ కురిసే అవకాశం ఉంది.

బలమైన ఈదురు గాలులు (గాలి వేగం గంటకు 30-40 కి.మీ.)తో తెలంగాణ రాష్ట్రంలోని అన్ని జిల్లాలలో అక్కడక్కడ వీచే అవకాశం ఉంది.

#### మొదవ రోజు:

ఉరుములు, మెరుపులతో కూడిన వర్షాలు తెలంగాణ రాష్ట్రంలోని ఆదిలాబాద్, కొమరంభీం ఆసిఫాబాద్, మంచిర్యాల, నిర్మల్, నిజామాబాద్, జగిత్యాల, రాజన్న సిరిసిల్ల, కరీంనగర్, పెద్దపల్లి, జయశంకర్

భూపాలపల్లి, ములుగు, భద్రాద్రి కొత్తగూడెం, ఖమ్మం, మహబూబాబాద్, వరంగల్, హన్మకొండ, వికారాబాద్, సంగారెడ్డి, మెదక్, కామారెడ్డి జిల్లాలలో అక్కడక్కడ కురిసే అవకాశం ఉంది.

బలమైన ఈదురుగాలులు (గాలి వేగం గంటకు 30-40 కి.మీ.)తో తెలంగాణ రాష్ట్రంలోని అన్ని జిల్లాలలో అక్కడక్కడ వీచే అవకాశం ఉంది.

**నాల్గవ రోజు:** బలమైన ఈదురుగాలులు (గాలి వేగం గంటకు 30-40 కి.మీ.)తో తెలంగాణ రాష్ట్రంలోని అన్ని జిల్లాలలో అక్కడక్కడ వీచే అవకాశం ఉంది.

**ఐదోవ రోజు:** బలమైన ఈదురుగాలులు (గాలి వేగం గంటకు 30-40 కి.మీ.)తో తెలంగాణ రాష్ట్రంలోని అన్ని జిల్లాలలో అక్కడక్కడ వీచే అవకాశం ఉంది.

### **వాతావరణ ఆధారిత వ్యవసాయ సలహాలు**

- భారీ వర్షాలు కురిసే అవకాశం ఉన్నందున పొలంలో నుండి మురుగు నీటిని తీసివేయడానికి కాలువలు చేసుకోవాలి.
- భారీ వర్షాలు కురిసిన ప్రాంతాలలో పొలం నుండి మురుగు నీటిని తీసివేయాలి.
- భారీ వర్షాలు కురిసే అవకాశం ఉన్నందున పంట పొలాల్లో మందులను పిచికారి చేయడం తాత్కాలికంగా వాయిదా వేయాలి.
- ఉరుములు మరియు మెరుపులతో కూడిన వర్ష సూచనలున్నందున రైతులు విద్యుత్ స్థంబాలు, విద్యుత్ తీగలు మరియు చెరువులు, నీటి కుంటలకు దూరంగా ఉండవలెను.
- రాష్ట్రంలో అక్కడక్కడ వివిధ జిల్లాలలో ఉరుములు మరియు మెరుపులతో కూడిన వర్ష సూచనలున్నందున రైతులు చెట్ల క్రింద నిలబడరాదు మరియు పశువులు, గొర్రెలు, మేకలను చెట్ల క్రింద ఉంచరాదు.
- ప్రస్తుతం కురిసిన వర్షాలను ఉపయోగించుకొని వర్షాధార పంటలకు పైపాటుగా నత్తజని మరియు పోటాష్ ఎరువులను సిఫారసు చేసిన మోతాదులో అందించాలి.
- రైతులు వర్షాధార పంటలైన ప్రత్తి, సోయాచిక్కుడు, మొక్కజొన్న, జొన్న, కంది, పెసర మొదలగు పంటలను నేల 15-20 సెం.మీ. లోతు తడిచిన తరువాతనే పంటలను విత్తుకోవాలి.
- వేరుశనగ, సోయాచిక్కుడు, జొన్న, పెసర, కంది మరియు మినుము పంటలను విత్తుకునే ముందు తప్పనిసరిగా విత్తన శుద్ధి చేసుకోవాలి.
- ఇప్పటివరకే ప్రత్తి విత్తుకొన్న రైతులు విత్తిన 7-10 రోజులలోపు మొలకరాని స్థానంలో మరల విత్తనం విత్తుకోవాలి.
- వరి తరువాత వరి సాగు చేసే రైతులు పచ్చిరొట్ట పైరును (జనుము, జిలుగ, పెసర, పిల్లి పెసర, ఉలవలు మొదలైన) విత్తుకోవడానికి అనుకూల సమయం.
- ముందస్తుగా వేసుకున్న పచ్చిరొట్ట పంటలను పూత దశలో భూమిలో కలియదున్నాలి.

### **వరి**

- జులై మాసం తెలంగాణ సోన (RNR 15048) వరి రకంతో నారుమడి పోయుటకు అనుకూలం.

- రైతులు మధ్యకాలిక (130-135 రోజుల) లేదా స్వల్పకాలిక (120-125 రోజుల) వరి రకాల నారుమళ్ళు జూన్ 20వ తేదీ నుండి జూలై 15వ తేదీ వరకు పోసుకోవచ్చును.
- ప్రస్తుతం 25-30 రోజుల వయస్సు ఉన్న దీర్ఘకాలిక వరి రకాలను నాటు పెట్టుకోవడాని సరైన సమయం.
- వరి నాట్లు వేసుకునే వారం రోజుల ముందు ఎకరాకు సరిపోయే నారుమడికి 800 గ్రా. కార్బోప్యూరాన్ 3సిజి గుళికలను ఇసుకలో కలిపి చలుకోవాలి.
- నాటు పెట్టిన తర్వాత ప్రతి 2మీటర్లకు కాళిబాటను తీయటం వలన గాలి వెలుతురూ బాగా ప్రసరించడంతో బాటు సుడిదోమ ఉద్యతిని నివారించవచ్చు. అదేవిధంగా రైతులు ఎరువులు మరియు పురుగు మందులు పంటకు అందించడానికి సులువుగా ఉంటుంది.

### ప్రత్తి

- ప్రత్తి పంటను జూలై 20వ తేదీ వరకు దిగుబడిలో ఎటువంటి తరుగుదల లేకుండా సాగు చేసుకోవచ్చు.
- ప్రత్తి వర్షాధార పంట కావున, రైతులు వారికి అనుకూలమైన బిటి రకాలను బోదెలు మరియు కాలువల పద్ధతిలో విత్తుకోవాలి.
- ఎక్కువ వర్షాలు పడినప్పుడు కూడా కాలువల ద్వారా నీటిని తీసివేయుటకు అనుకూలం.
- ప్రస్తుతం కురిసిన వర్షాలను ఉపయోగించుకొని వర్షాధార ప్రత్తి పంటలో మొదటి దఫా మరియు రెండవ దఫా పైపాటు నత్తజని మరియు పొటాషియం ఇచ్చే ఎరువులను 20 మరియు 40 రోజుల అందించాలి.

### మొక్కజొన్న

- మొక్కజొన్న పంటను జూలై 15వ తేదీ వరకు మధ్య మరియు దీర్ఘకాలిక (90-100 రోజులు మరియు 100-120 రోజులు) రకాలను దిగుబడిలో ఎటువంటి తరుగుదల లేకుండా విత్తుకొనుటకు అనుకూల సమయం.
- మొక్కజొన్న శాకీయ దశలో ఎక్కువ నీటిని మరియు ప్రత్యుత్పత్తి దశలో నీటి ఎద్దడిని తట్టుకోలేదు కాబట్టి పంటను బోదెలు మరియు కాలువల పద్ధతిలో సాగు చేసుకోవలెను.
- 14రోజుల వయస్సు గల మొక్కజొన్న పంటలో కత్తిర పురుగు నివారణకు ముందు నివారణ చర్యలలో భాగంగా వేప నూనెను పిచికారి చేసుకోవాలి.

### ఆముదం

- ఆముదం పంటను జూలై రెండవ పక్షం వరకు విత్తుకోవచ్చును.

### కంది

- కంది పంటను జూలై 15వ తేదీ వరకు దిగుబడిలో ఎటువంటి తరుగుదల లేకుండా సాగు చేసుకోవచ్చు.
- కందిని ఎకపంటగా లేదా ప్రత్తి, మొక్కజొన్న, జొన్న, సోయాచిక్కుడు, పెసలు మరియు మినుములతో అంతర పంటగా సాగు చేసుకోవచ్చు.
- కంది పంటను విత్తే ముందు విత్తనాని 2.5గ్రా. కాప్టాన్ లేదా 3గ్రా. థైరామ్ మరియు 10గ్రా. ట్రికోడేర్మా విరిడె మందుతో కిలో విత్తనానికి కలిపి విత్తుకున్నట్లయితే విత్తనం లేదా భూమి ద్వారా సంక్రమించే శీలీంధ్రాల నుండి పంటను కాపాడవచ్చు.

### పెసర మరియు మినుము

- పెసర మరియు మినుము పంటను జులై 15వ తేదీ వరకు విత్తుకోవచ్చును.
- పెసర మరియు మినుమును ఎకపంటగా లేదా ప్రత్తి మరియు కందితో అంతర పంటగా సాగు చేసుకోవచ్చు.
- వరి సాగు చేసే పొలాల్లో, ఆలస్యంగా నీరు విడుదలయ్యే ప్రాంతాలలో పెసరను పైరుగా లేనిచో పచ్చి రొట్టగా విత్తుకోవాలి.

### సోయాచిక్కుడు

- సోయాచిక్కుడు పంటను జులై మొదటి వారం వరకు విత్తుకోనుటకు అనుకూల సమయం.
- నీటి వసతి గల తేలికపాటి నేలలు కూడా సోయాచిక్కుడు సాగుకు అనుకూలం.
- ఎత్తు మడులు మరియు కాలువల (Broad Bed and Furrow – BBF) పద్ధతిలో విత్తుకున్నట్లయితే విత్తన మోతాదు తగ్గటంతోపాటు సరైన సాంద్రతలో మొక్కలు ఉండి, సాగు ఖర్చులు తగ్గి అధిక దిగుబడులు వస్తాయి.
- ఈ పద్ధతిలో భూమిలో ఎక్కువ తేమ నిలువ ఉండి పంట నీటి యద్దడిని తట్టుకుంటుంది. అదేవిధంగా ఎక్కువ నీటిని పొలం నుండి తీసివేయటానికి కాలువలు ఉపయోగపడతాయి.

### చెరకు

- తోట వయస్సు నాలుగు నెలలప్పుడు (జూన్ – జులై మాసాల్లో) మొక్కల వరుసల మొదళ్ళకు ఎత్తుగా మట్టిని ఎగదోయాలి. దీని వలన అధిక వర్షాలు మరియు తుఫాను గాలులకు పంట పడిపోకుండా కాపాడుకోవచ్చు.

### కూరగాయలు

- తీగ జాతి కూరగాయ పంటలు విత్తుటకు అనుకూలం.
- మిరప, టమాట, వంగ మరియు బంతి పంటలను ఎత్తు మడులమీద నారు పోయడం ద్వారా నారుకుళ్ళు తెగులు నివారించవచ్చు.

### మామిడి

- రైతులు కోత అనంతరం 2-4 వారాల తరువాత ఎండు కొమ్మలను, తెగులు సోకిన కొమ్మలను మరియు అడ్డంగా పెరిగిన కొమ్మలను కత్తిరించినట్లయితే గాలి మరియు వెలుతురు మొక్కకు బాగా లభిస్తుంది.
- తోటలో రాలిపోయిన, కుళ్ళిపోయిన, తెగుళ్ళు ఆశించిన కాయలతో పాటు ఎండుఆకులను మరియు కొమ్మలను సేకరించి తోటలకు దూరంగా కాల్చివేయాలి.

### పశుపోషణ

- ప్రస్తుత వాతావరణ పరిస్థితులు కోళ్ళలో కొక్కిర తెగులు సోకుటకు అనుకూలం వీటి నివారణకు, టీకాలు వేయించాలి.
  - గొర్రెలలో చిటుకు మరియు PPR
  - ఆవులు మరియు గేదెలలో గొంతువాపు వ్యాధి సోకుటకు అనుకూలం వీటి నివారణకు, టీకాలు వేయించాలి.
  - గొర్రెలలో నట్టల నివారణకు డీవార్మింగ్ చేయవలెను.

# PROFESSOR JAYASHANKAR TELANGANA AGRICULTURAL UNIVERSITY

Agro Climate Research Unit (ACRU),  
Agricultural Research Institute, Rajendranagar, Hyderabad-30.



Bulletin No. XXXVI/54/2025

## WEATHER BASED AGROMET ADVISORIES FOR TELANGANA STATE

FROM DATE: 09.07.2025 (Wednesday) TO 13.07.2025 (Sunday)



Grameen Krishi Mausam Seva (GKMS Project)

INDIA METEOROLOGICAL DEPARTMENT, NEW DELHI.

### Weather details of past 3 days:

During last three days, light to moderate rains occurred at many places over different parts of the State. The maximum and minimum temperatures ranged between 29°C to 35°C and 20°C to 26°C respectively.

### Weather forecast for the next 5 days (Based on today's forecast at 1300 hrs):

As per the forecast received from Meteorological Centre, Hyderabad, light to moderate rains or thundershowers very likely to occur at few places over different parts of the state during next five days. The maximum and minimum temperatures are likely to range between 26°C to 36°C and 21°C to 28°C respectively.

### Warning:

**Day 1:** Heavy to Very Heavy Rain accompanied with Thunderstorm, Lightning and Gusty winds (40-50 kmph) very likely to occur at isolated places in Adilabad, Kumarambheem Asifabad, Nirmal districts of Telangana.

Heavy Rain accompanied with Thunderstorm, Lightning and Gusty winds (30- 40 kmph) very likely to occur at isolated places in Mancherial, Nizamabad districts of Telangana.

Thunderstorm accompanied with Lightning very likely to occur at isolated places in Jagtial, Rajanna Sirisilla, Karimnagar, Peddapalli, Jayashankar Bhupalpally, Mulugu, Bhadradi Kothagudem, Khammam, Nalgonda, Suryapet, Mahabubabad, Warangal, Hanamkonda, Jangaon, Siddipet, Yadadri Bhuvanagiri, Rangareddy, Hyderabad, Medchal Malkajgiri, Vikarabad, Sangareddy, Medak, Kamareddy districts of Telangana.

Strong surface winds up to (30-40 kmph) very likely to prevail at isolated places in ALL districts of Telangana.

**Day 2:** Heavy Rain accompanied with Thunderstorm, Lightning and Gusty winds (30- 40 kmph) very likely to occur at isolated places in Adilabad, Kumarambheem Asifabad, Mancherial, Nirmal districts of Telangana.

Thunderstorm accompanied with Lightning very likely to occur at isolated places in Nizamabad, Jagtial, Rajanna Sirisilla, Karimnagar, Peddapalli, Jayashankar Bhupalpally, Mulugu, Bhadradi Kothagudem, Khammam, Nalgonda, Suryapet, Mahabubabad, Rangareddy, Hyderabad, Medchal Malkajgiri, Vikarabad, Sangareddy, Medak, Kamareddy districts of Telangana.

Strong surface winds up to (30-40 kmph) very likely to prevail at isolated places in ALL districts of Telangana.

**Day 3:** Thunderstorm accompanied with Lightning very likely to occur at isolated places in Adilabad, Kumarambheem Asifabad, Mancherial, Nirmal, Nizamabad, Jagtial, Rajanna Sirisilla, Karimnagar, Peddapalli, Jayashankar Bhupalpally, Mulugu, Bhadradi Kothagudem, Khammam, Mahabubabad, Warangal, Hanamkonda, Vikarabad, Sangareddy, Medak, Kamareddy districts of Telangana.

Strong surface winds up to (30-40 kmph) very likely to prevail at isolated places in ALL districts of Telangana.

**Day 4:** Strong surface winds up to (30-40 kmph) very likely to prevail at isolated places in ALL districts of Telangana.

**Day 5:** Strong surface winds up to (30-40 kmph) very likely to prevail at isolated places in ALL districts of Telangana.



## **Weather based Agro Advisories**

- In view of heavy rainfall forecast, provide drainage channels to drain out excess water from the crop fields.
- Drain out excess water from the fields in areas received heavy rainfall.
- In view of heavy rainfall forecast, temporarily postpone the spraying of chemicals in the agricultural fields.
- In view of the thunderstorm accompanied with lightning forecast, farmers are advised not to stand under the trees and don't allow the Cattle / Goats / Sheep's to take shelter under trees.
- In view of the thunderstorm accompanied with lightning forecast, move away from water bodies, electric poles and wires.
- In view of the receipt of the rainfall advised to go for top dressing of Nitrogen and Potassic fertilizers for rainfed crops.
- Farmers are advised to go for sowing of rainfed crops (Cotton, Soybean, Maize, Jowar, Redgram, Greengram, Cotton etc.,) after the soil wet up to top 15-20 cm depth.
- Treat the seeds of Groundnut, Soybean, Jowar, Greengram, Redgram and Blackgram before sowing.
- Farmers who have already sown cotton crop are advised to go for gap filling within 7-10 days after sowing.
- Rice followed by rice cropping system farmers are advised to go for sowing of *in-situe* green manure crops (Dhaincha, Sunhemp, Greengram, Pillipesara, Horsegram etc.,).
- Green manure crops at flowering stage are advised to incorporate into soil.

### **Rice**

- Advised to take up the paddy nurseries using Telangana Sona (RNR 15048) in the month of July.
- Raising of rice nurseries of medium duration (130-135 days) or short duration (120-125 days) varieties is under progress and continuous until 15<sup>th</sup> July.
- It is time to transplant long duration varieties with 25-30 days old seedlings.
- Farmers are advised to apply the Carbofuran 3CG granules @ 800 g per acre of nursery one week before pulling of the nursery.
- After transplanting form alleyways at 2mtrs interval to allow the free flow of the air and other natural resources, to control BPH and also help the farmer to take up the other management practices like fertilizer and pesticide application.

### **Cotton**

- Sowings is under progress and continuous until July 20<sup>th</sup> without any yield reduction
- Sow the crop on ridge and furrow method to conserve the moisture by selecting suitable BT varieties as this method helps to remove the excess water from the fields.
- Top dressing of first dose of Nitrogen and Potassic fertilizers is recommended at 20 DAS whenever there is soil moisture.

### **Maize**

- Sowings is under progress and continuous until July 15<sup>th</sup> with medium and long duration varieties (90-100 days and 100-120 days) without any yield reduction.
- Sow the crop in ridge and furrow method to with stand excess moisture stress during the vegetative stage and less moisture stress at tasseling and silking stage.
- Initiate the Fall Army Worm control measures 14 days after sowing, with neem oil as a preventive measure.

### **Castor**

- Sowings is under progress and continuous until July 15<sup>th</sup>.



### **Redgram**

- Sowings is under progress and continuous until July 15<sup>th</sup> without any yield reduction.
- Crop can be sown either as sole crop or as intercrop with Cotton, Maize, Jowar, Soybean, Greengram and Blackgram.
- Seed should be treated with Captan @ 2.5g or Thiram @ 3g + Trichoderma Viride @ 10g per kg seed to protect the crop from seed or soil borne fungi.

### **Greengram and Blackgram**

- Sowings is under progress and continuous until July 15<sup>th</sup>.
- Crop can be sown either as sole crop or intercrop with Cotton and Redgram.
- Take the Greengram as catch crop to preceding rice in areas where the release of water is delayed.

### **Soybean**

- Sowings is under progress and continuous until July 1<sup>st</sup> week without any yield reduction.
- Light soils with assured irrigation are also suitable for soybean cultivation.
- Use Broad Bed Furrow (BBF) method of sowing to reduce seed rate, cost of sowing, with uniform germination and finally to get higher yields.
- By this method the crop can with stand moisture stress and at the same time furrows are useful to remove excess water from the fields.

### **Sugarcane**

- In four months old crop (June – July) earthing up should be carried out. This will protect the crop from lodging due to heavy rains and stormy/gusty winds.

### **Vegetables**

- Take up sowings of Gourds (Ridge Gourd / Snake Gourd / Bitter Gourd etc.,).
- Take up the nursery of Chillies, Tomato, Brinjal and Marigold on raised beds to avoid damping off.

### **Mango**

- Farmers are advised to go for pruning of dried, diseased and crisscross branches 2-4 weeks after final harvest to allow the proper ventilation and distribution of solar radiation.
- Advised to collect the rotten fruits and fallen leaves and burn them outside the mango orchard.

### **Livestock**

- Prevailing weather conditions are congenial for the occurrence of
  - Ranikhet disease in poultry, ET and PPR in Sheep
  - Hemorrhagic Septicemia, Foot and mouth disease in cattle and Buffaloes. To prevent the diseases, vaccinate the animals
  - Deworming in Sheep

Principal Scientist (Agro) & Head